

Table 1: A comparison of the range in median PFAS levels detected in the PIVUS cohort to other background level general trend studies.

PFAS	PIVUS (2001-2014)	The U.S.(1) (2003-2012)	Germany(2) (2001-2010)	Australia(3) (2002-2011)	Norway(4) (2001-2007)
Range of median concentrations (ng/mL)					
PFHpA	0.03-0.07	<LOD	N/A	N/A	0.1-0.1
PFOA	2.5-4.0	2.16-4.3	3.2-5.2	4.3-9.7	3.1-4.2
L-PFOS	7.6-14	N/A	N/A	N/A	23-30
PFNA	0.73-0.87	0.89-1.23	N/A	N/A	1.1-1.5
PFDA	0.32-0.34	<LOD-0.300	N/A	N/A	0.7-0.8
PFUnDA	0.30-0.45	<LOD-0.200	N/A	N/A	1.1-1.3
FOSA	0.02-0.11	<LOD-0.100	N/A	N/A	0.2-1.0
PFHxS	2.2-3.5	1.28-1.90	0.86-1.87	N/A	1.9-2.0




Figure 1: A comparison of the longitudinal change in concentration of four PFASs detected in the PIVUS cohort to the results obtained from cross-sectional and retrospective trend studies conducted in the U.S. (1), Australia (3), Germany (2) and Norway (4).