

Cod: 8.4007

## HIGHLIGHTS FROM THE NEW AMAP ASSESSMENT REPORT ON CHEMICALS OF EMERGING ARCTIC CONCERN

C.A. De Wit<sup>1</sup>, J. Balmer<sup>2</sup>, D. Muir<sup>3</sup>, K. Vorkamp<sup>4</sup>, S. Wilson<sup>2</sup>

<sup>1</sup>*Department of Environmental Science and Analytical Chemistry (ACES), Stockholm University, Sweden*

<sup>2</sup>*Arctic Monitoring and Assessment Programme, Oslo, Norway*

<sup>3</sup>*Canada Centre for Inland Waters, Environment and Climate Change Canada*

<sup>4</sup>*Department of Environmental Science, Aarhus University, Denmark*

### Introduction

The Arctic Monitoring and Assessment Programme (AMAP) previously produced three assessment reports on persistent organic pollutants (POPs) in the Arctic<sup>1-3</sup>. A new fourth update assessment<sup>4</sup> addresses ‘chemicals of emerging Arctic concern’, including chemicals with POP-like characteristics and also some chemicals and groups of substances that may not meet the classical definition of POPs. Under the Stockholm Convention, one of the criteria for establishing that chemicals are persistent in the environment is if they are found at locations “distant from sources” or “where monitoring data show that long-range environmental transport of the chemical...may have occurred”<sup>5</sup>. The new assessment compiles available Arctic data for per- and polyfluorinated substances (PFAS), brominated, chlorinated and organophosphorus-based flame retardants and plasticisers (BFRs, CFRs, PFRs), phthalates, short chain chlorinated paraffins (SCCPs), siloxanes, pharmaceuticals and personal care products (PPCPs), polychlorinated naphthalenes (PCNs), hexachlorobutadiene, current use pesticides (CUPs), pentachlorophenol/ pentachloroanisole, non-Aroclor and byproduct PCBs, halogenated natural products (HNPs), mono- and dibutyltins, polycyclic aromatic hydrocarbons (PAHs), and microplastics.

### Methods

The assessment was based on a review of both the recent peer reviewed and grey literature for existing data on a wide range of chemicals of emerging concern (CECs) that have recently been detected in Arctic environmental compartments. The emphasis was on environmental levels and trends.

### Results and discussion

Results from air monitoring at Arctic sites (including Alert, Canada; Pallas, Finland; Zeppelin station, Svalbard; Storhofdi, Iceland; Nord, Greenland and shipboard measurements on Arctic expeditions) confirm that, like ‘traditional’ POPs (e.g. PCBs and DDT), many CECs also undergo long-range atmospheric transport from source regions to the Arctic. These include PFAS, BFRs, CFRs, PFRs, SCCPs, phthalates, siloxanes, PCNs, HCBd, several CUPs, and PAHs. Results for several PFRs in air showed surprisingly high concentrations compared to sum-PBDEs. At least seven new pesticides were measured in Arctic media: MCPA, metribuzin, pendimethalin, phosalone, quizalofop-ethyl, tefluthrin and triallate.

Studies of biota in the Arctic terrestrial environment are few, but show that some CECs are found in reindeer. In Arctic freshwater and marine environments, a number of CECs (long chain PFCAs, several new BFRs, dechloranes, PFRs, phthalates, SCCPs, siloxanes, some PPCPs, PCNs, HCBd, several CUPs, organotins, some HNPs) have been found in fish, seabirds and marine mammals, indicating bioaccumulation potential for these. For some of these, (long chain PFCAs, a few BFRs ( $\alpha$ -HBCDD, DBDPE), SCCPs, and PCNs) biomagnification was also indicated. Figure 1 illustrates the trophic magnification of SCCPs in an Arctic food web from Svalbard.

Sparse data mean that few spatial trends were discernible, and only for Canada, Greenland, Svalbard and the European Arctic, with highest concentrations of many CECs found on Greenland and Svalbard. Some temporal trends were available in air, indicating stable or increasing concentrations of some new BFRs, SCCPs, some CUPs, and for biota, long chain PFCAs. An example of temporal trends for several CUPs is shown in Figure 2. However, more data for most CECs will be required for a more comprehensive assessment of trends.

For some CECs, concentrations were higher in proximity to settlements and urban sites. This was particularly apparent for several CECs that are used in consumer products. In particular, siloxanes and PPCPs were found locally in receiving waters impacted by (often untreated) sewage effluents from Arctic communities. Their persistence is enhanced due to cold conditions and periods of no light, which slow down microbial- and photo-degradation. Thus, several CECs (including PPCPs, siloxanes, phthalates) differ from traditional POPs in that localized, point sources of pollutants within the Arctic may be important. However, the importance of local sources compared to long-range transport needs to be assessed.

Indications of long-range transport, persistence and bioaccumulation potential indicate that some CECs, such as long chain PFCAs, PCNs, SCCPs and some BFR/CFRs may be possible POPs as defined by the Stockholm Convention, and constitute an 'emerging concern' with respect to contamination of the Arctic.

Many CECs have been analyzed using previously established methods for other, similar classes of persistent chemicals. For some of the emerging chemicals, cleanup and analytical methods have only recently been developed and these are not yet standardized. QA/QC procedures for new contaminants are not always optimal as pure standards may not be commercially available yet. For most of the CECs, there are no standard reference materials with certified concentrations available to validate methods. Because many emerging chemicals are used in consumer products and building materials, there are also considerable risks with contamination of low level samples. Thus, there is a need for interlaboratory comparison studies, development of best practice for analysis, increase in the number of commercially available standards and certified reference materials for many of the CECs addressed in this assessment.

#### **Acknowledgements**

We thank the many authors and contributors to the assessment report as well as national funding authorities for making their participation possible.

#### **References**

1. de March, B. G. E., de Wit, C.A., Muir, D.C.G., Braune, B.M., Gregor, D.J., Norstrom, R.J., Olsson, M., Skaare, J.U., and Stange, K. (1998) In AMAP Assessment Report. Arctic Pollution Issues, pp. 183-372, Arctic Monitoring and Assessment Programme, Oslo, Norway.
2. de Wit, C. A., Fisk, A. T., Hobbs, K. E., Muir, D. C. G., Gabrielsen, G. W., Kallenborn, R., Krahn, M. M., Norstrom, R. J., and Skaare, J. U (2004) AMAP Assessment 2002: Persistent Organic Pollutants in the Arctic, xvi +310 pp, Arctic Monitoring and Assessment Programme, Oslo, Norway.
3. AMAP (2010) Science of the Total Environment Special Issue, 408, 2851-3051.
4. AMAP (to be published 2016) Chemicals of Emerging Arctic Concern, see [www.amap.no](http://www.amap.no).
5. UNEP (2009) Stockholm Convention on Persistent Organic Pollutants (POPs) as amended in 2009. United Nations Environment Programme, Geneva, CH. (<http://chm.pops.int/TheConvention/Overview/TextoftheConvention/tabid/2232/>)
6. NILU (2013) Rapport: M40-2013. Norwegian Institute for Air Research (NILU) Oslo, Norway, 110 pp.
7. Jantunen, L. (2014) Synopsis of Research Conducted under the 2013–2014 Northern Contaminants Program. Ottawa ON, Aboriginal Affairs and Northern Development Canada: pp 329-339.
8. Jantunen, L. M., Wong, F., Gawor, A., Kylin, H., Helm, P.A., Stern, G.A., Strachan, W.M.J., Burniston, D.A. and Bidleman, T.F. (2015) Environmental Science and Technology 49, 23, 13844–13852.




Figure 1. Relationship between SCCP concentrations (log ng/g lw) and  $\delta^{15}\text{N}$  in a food web from Svalbard. Figure reproduced from NILU<sup>6</sup>.




Figure 2. Temporal trends of current use pesticides (CUPs) measured in air of the western Arctic Ocean collected during oceanographic expeditions held between 1999-2013. Arithmetic mean concentrations are given in  $\text{pg}/\text{m}^3$ . Figure from Jantunen<sup>7</sup>; data listed in Jantunen et al.<sup>8</sup> Expeditions, listed by year, traversed the Bering-Chukchi Seas (BERPAC-1993), central Canada Basin to the North Pole (AOS-1994) and the Canadian Archipelago (TNW-1999, IPY-2007, IPY-2008, ArcticNet-2010, ArcticNet-2011, ArcticNet-2013).