





In conclusion, this study confirmed that PFOA accumulates in bone also in humans, is eliminated very slowly and can affect bone homeostasis both directly via bone cells, and indirectly via increased body weight and decreased activity. PFOA levels were also detectable in control mice, suggesting that even in controlled laboratory environment exposure is possible.

### **Acknowledgements**

We thank Miia Vierimaa and Arja Moilanen for technical assistance and FORMAS for support.

### **References**

- Bogdanska J, Borg D, Sundström M et al. *Toxicology* 2011;284:54-62.  
Chan BY, Gartland A, Wilson PJ et al. *Bone* 2007;40:149-159.  
ECHA, Proposals to identify Substances of Very High Concern current consultations - ECHA. 2013.  
Giaginis C, Tsantili-Kakoulidou A, Theocharis S. *Fundam Clin Pharmacol* 2007;21:231-244.  
Korkalainen M, Kallio E, Olkku A. et al. *Bone* 2009;44:1134-1142.  
Koskela A, Tuukkanen J, Viluksela M et al. *Toxicol Appl Pharmacol* 2012;263:210-217.  
Maloney EK, Waxman DJ. *Toxicol Appl Pharmacol* 1999;161:209-218.  
Perez F, Nadal M, Navarro-Ortega A et al. *Environ Int* 2013;59:354–362.  
Tsukamoto Y, Ishihara Y, Miyagawa-Tomita S et al. *Biochem Pharmacol* 2004;68:739-746.  
Zareitalabad P, Siemens J, Hamer M, et al. *Chemosphere* 2013;91:725-732.