


Figure 1 Distributions of Σ 3HBCDs in surface waters and sediments in Weifang City, Shandong Province, East China



References:

1. Yi S, Liu J G, Jin J, et al. Assessment of the occupational and environmental risks of hexabromocyclododecane (HBCD) in China. *Chemosphere*, 2016, 150:431-7.
2. LETCHER R J, LU Z, CHU S, et al. Hexabromocyclododecane Flame Retardant Isomers in Sediments from Detroit River and Lake Erie of the Laurentian Great Lakes of North America. *Bulletin of Environmental Contamination and Toxicology*, 2015, 95: 31-6.
3. De Wit C A, Herzke D, Vorkamp K. Brominated flame retardants in the Arctic environment - trends and new candidates. *Science of the Total Environment*, 2010, 408(15): 2885-918.
4. Zhu H, Zhang K, Sun H, et al. Spatial and temporal distributions of hexabromocyclododecanes in the vicinity of an expanded polystyrene material manufacturing plant in Tianjin, China. *Environmental Pollution*, 2017, 222(338-47).
5. Anim A K, Drage D S, Goonetilleke A, et al. Distribution of PBDEs, HBCDs and PCBs in the Brisbane River estuary sediment. *Marine pollution bulletin*, 2017, 120(1-2): 165-73.
6. Wang L, Zhang M, Lou Y, et al. Levels and distribution of tris-(2,3-dibromopropyl) isocyanurate and hexabromocyclododecanes in surface sediments from the Yellow River Delta wetland of China. *Marine Pollution Bulletin*, 2017, 114(1): 577-82.