

SEASONAL VARIATION OF PCDD/PCDF AND dl-PCB CONCENTRATIONS IN AMBIENT AIR USING PASSIVE SAMPLERS

Sau TK¹, Truong NX¹, Hung LB¹, Thang ND¹, Trien DH¹, Nam PD, Long NP¹, Long LV¹

¹Vietnam-Russia Tropical Centre, Nguyen Van Huyen Str., Hanoi, Vietnam, 100000, sau_tk@yahoo.com

1. INTRODUCTION

Monitoring of PCDDs/PCDFs and dl-PCBs in ambient air using passive air sampler (PAS) with polyurethane foam (PUF) disk at sampling intervals of approximately three months was conducted in Hanoi and Danang, Vietnam¹⁻⁴. The ¹³C-labeled PCDD/PCDF standards were spiked on the PUF disks as surrogates at the beginning of each sampling interval. The seasonal variation of PCDD/PCDF and dl-PCB concentrations was studied by Spring, Summer, Autumn and Winter in 7 continuous years from 2012 to 2018 in Hanoi. This paper presents the results of seasonal monitoring and the temporal trends of PCDDs/PCDFs and dl-PCBs in ambient air between Winter 2015 - Autumn 2018 and in comparison with our previous study between Spring 2012 - Autumn 2015⁵.

2. MATERIALS AND METHODS

2.1. Passive air sampler

The TE-200 PAS sampler of Tisch Environmental Inc (Ohio, USA) with the PUF disk was hanged outdoor⁶. The PUF disks were pre-cleaned by Soxhlet extraction with acetone, then with toluene and were dried in a vacuum oven, then stored in an air-tight container before use.

2.2. Standards of PCDDs/PCDFs and dl-PCBs

The surrogate standards consisted of 15 isotopes of ¹³C-labeled PCDDs/PCDFs diluted from the stock solution EDF-8999 of Cambridge Isotope Laboratories (Massachusetts, USA). The native PCDD/PCDF standards, and other ¹³C-labeled recovery and ³⁷Cl-labeled clean-up standards were used as described in the US EPA method 1613B⁷. The native and ¹³C-labeled dl-PCB standards were used according to the US EPA method 1668B⁸.

2.3. Experiments

The experiments were conducted seasonally at an urban area in Hanoi and were described as in the previous study⁵. The PAS sample IDs and sampling intervals are shown in Table 1. The sample IDs with the additional suffix (-D) were the field duplicate PAS samples. Their UTM coordinates were 582988E-2327289N and 582984E-2327299N, respectively.

Table 1: PAS sample IDs and sampling intervals

Season	Spring	Summer	Autumn	Winter
2015				HNP33, HNP34-D (03 Nov - 03 Feb 2016)
2016	HNP35, HNP36-D (03 Feb - 04 May)	HNP37, HNP38-D (04 May - 03 Aug)	HNP39, HNP40-D (03 Aug - 02 Nov)	HNP41, HNP42-D (02 Nov - 07 Feb 2017)
2017	HNP43, HNP44-D (07 Feb - 05 May)	HNP45, HNP46-D (05 May - 14 Aug)	HNP48 (14 Aug - 09 Nov)	HNP50 (14 Nov - 26 Feb 2018)
2018	HNP53 (26 Feb -18 May)	HNP56 (18 May - 15 Aug)	HNP59 (15 Aug - 14 Nov)	

2.4. Analysis of PCDDs/PCDFs and dl-PCBs

The recovery standards of ¹³C-1234-TCDD and ¹³C-123789-HxCDD were added before PAS sample extraction to determine the efficiency of surrogate retention on the passive PUF disks. Then PAS samples were prepared and analyzed as described in the US EPA method 1613B⁷ and method 1668B⁸.

PCDDs/PCDFs and dl-PCBs were analyzed by high resolution gas chromatography (HRGC Aligent 7890A) and high resolution mass spectrometry (HRMS AutoSpec Premier, Water). Temperature program: 150°C for 2 minutes, increasing from 150°C to 220°C at 20°C/min, 220°C for 16 minutes, from 220°C to 320°C at 5°C/min and keeping at 320°C until finish. Temperature of the injector: 280°C, interface: 290°C. Helium: 1.0 ml/min. DB-5MS column: 60 m length, 0.25 mm id., 0.25 µm film thickness The MS was operated at resolution $\geq 10,000$ at 10% valley under positive electron ionization.

LODs of method are 0.2 pg/sample for TetraCDDs/CDFs, 0.5 pg/sample for Pen- through Hepta-CDDs/CDFs, and 1.0 pg/sample for OCDD/OCDF and dl-PCBs.




Figure 4: Concentration of PCDDs/PCDFs, dl-PCBs and their temporal trend in Autumn

The figures 1,2,3,4 and tables 2,3 show that the concentration of PCDDs/PCDFs and Total TEQ_{D/F} between Winter 2015 and Autumn 2018 were lowest in summers and highest in autumns. The variation of total TEQ_{D/F} (pg/disk/day) was from 0.196 to 0.351 (average 0.255) in summer; 0.334-0.371 (average 0.351) in autumn; 0.241-0.378 (0.309) in winter; and 0.282-0.352 (0.302) in spring. The concentrations of total PCDDs/PCDFs (pg/disk/day) were in the range as follows: 2.12-3.39 (2.56) in summer; 3.18-3.41 (3.33) in autumn; 2.34-3.62 (2.96) in winter; and 2.45-3.39 (2.70) in spring. In all seasons, almost PCDF congeners were predominant. The concentrations of total PCDFs in ambient air were always 1.7- to 2.6-fold higher than total PCDDs. The principal PCDD congeners were 1234678-HpCDD and OCDD, in there the concentrations of OCDD were 1.6- to 2.4-times greater than 1234678-HpCDD. The average concentration of the most toxic congener 2378-TCDD contributed from 8.3% to 17.2% to the total TEQ_{D/F} and from 7.2% to 14.9% to the total TEQ_{D/F&dl-PCB}.

The concentrations of Total dl-PCBs between Winter 2015 and Autumn 2018 were lowest in winters and highest in summers (Table 2). The major dl-PCB congeners in all seasons were PCB#118, then PCB#105 and PCB#77 (Figures 1,2,3 and 4). PCB#118 was always predominant in profiles of dl-PCB congeners. The average seasonal concentration of PCB#118 contributed from 52.1% to 69.9% to Total dl-PCBs. The variation of Total dl-PCB concentrations (pg/disk/day) was in the range of 20.9-36.9 (average 27.0) in winters; 24.5-48.4 (average 33.0) in springs; 38.5-104 (average 75.6) in summers; and 43.0-80.0 (average 55.6) in autumns. The seasonal average of Total TEQ_{dl-PCB} was only in the range of 0.034-0.053 pg/disk/day (Table 3) and insignificantly contributed from 10.1% to 14.1% to the total TEQ_{D/F&dl-PCB}.

Table 2: Seasonal variation of PCDD/PCDF and dl-PCB concentrations (pg/disk/day) in the ambient air between 2015 and 2018

Season	Total PCDDs			Total PCDFs			Total dl-PCBs		
	Min	Average	Max	Min	Average	Max	Min	Average	Max
Winter	0.648	1.04	1.42	1.59	1.92	2.21	20.9	27.0	36.9
Spring	0.876	0.985	1.06	1.48	1.72	2.35	24.5	33.0	48.4
Summer	0.658	0.771	0.875	1.47	1.79	2.64	38.5	75.6	104
Autumn	0.909	0.931	0.958	2.27	2.40	2.48	43.0	55.6	80.0

Table 3: Seasonal variation of Total TEQ_{D/F}, Total TEQ_{dl-PCB} and their Total TEQ (pg/disk/day) in the ambient air between 2015 and 2018

Season	Total TEQ _{D/F}			Total TEQ _{dl-PCB}			Total TEQ _{D/F&dl-PCB}		
	Min	Average	Max	Min	Average	Max	Min	Average	Max
Winter	0.241	0.309	0.378	0.037	0.051	0.063	0.285	0.360	0.424
Spring	0.282	0.302	0.352	0.025	0.034	0.038	0.320	0.336	0.377
Summer	0.196	0.255	0.351	0.021	0.038	0.048	0.239	0.293	0.372
Autumn	0.334	0.351	0.371	0.050	0.053	0.061	0.384	0.404	0.421

The assessment of variations and temporal trends of PCDD/PCDF concentrations in ambient air through two phases from 2012 to 2018 are illustrated in Figure 5. The concentrations of PCDDs/PCDFs in four seasons between Spring 2012 and Autumn 2015 were presented in our previous study⁵.

Generally, the concentrations of Total PCDDs in all seasons between Winter 2015 and Autumn 2018 were 1.03- to 1.5-fold lower than that in comparison with our previous study between Spring 2012 and Autumn 2015⁵. The seasonal concentrations of Total PCDFs and Total TEQ_{D/F} between Winter 2015 and Autumn 2018 were also 1.04- to 1.5-fold lower in winters and autumns, however, were approximately 1.1-fold higher in spring and summer between Spring 2012 and Autumn 2015. Especially, the variation of PCDD/PCDF concentrations in ambient air between Winter 2015 and Autumn 2018 was shown in a narrower range compared to the period of Spring 2012 and Autumn 2015 (Figure 5).




Figure 5: Seasonal variation of PCDD/PCDF concentrations (pg/disk/day) in the ambient air between 2012-2015 and 2015-2018

The relative percent difference (RPD) between PCDD/PCDF and dl-PCB concentrations of the field duplicate PAS samples was calculated as follows: $RPD\% = [(C1 - C2) / (C1 + C2) / 2] * 100\%$ ⁹. In which: C1 represents the PCDD/PCDF concentration on the PUF disk with spiking ¹³C-labeled PCDD/PCDF surrogates; and C2 represents the PCDD/PCDF concentration on the PUF disk without spiking surrogates. Approximately 87% of PCDD/PCDF concentrations and 79% of dl-PCB concentrations had the RPD < 40%. This agreement of PCDD/PCDF and dl-PCB concentrations from the field duplicate PAS samples in this study shows that the seasonal monitoring of PCDDs/PCDFs and dl-PCBs using PAS is reliable and suitable in the monsoon tropical climatic conditions in Northern Vietnam.

Acknowledgements

This work was supported by Vietnam-Russia Tropical Centre. We gratefully thank the Vietnam-Russia Tropical Centre for their assistance and encouragement in carrying on this study.

References:

1. Sau TK, Thu NT, Dung NT (2013) *Organohalogen Compounds*. 75: 418-421.
2. Sau TK, Truong NX (2013) *Organohalogen Compounds*. 75: 422-425.
3. Sau TK, Truong NX, Hung LB, et al. (2014) *Organohalogen Compounds*. 76: 98-101.
4. Sau TK, Truong NX (2014) *Organohalogen Compounds*. 76: 101-105.
5. Sau TK, Truong NX (2016) *Organohalogen Compounds*. 78: 1068-1072.
6. Harner T (2006) PUF Disk Sampler (TE-200 PAS) Standard Operating Procedure.
7. US EPA (1994) *Method 1613B: Tetra- through octa-chlorinated dioxins and furans by isotope dilution HRGC/HRMS*.
8. US EPA (2008) *Method 1668B: Chlorinated biphenyl congeners in water, soil, sediment, biosolids, and tissue by HRGC/HRMS*.
9. US EPA (2014) *Hazardous waste test methods SW-846*. Chapter one: Project quality assurance and quality control.