

Figure 1. Elimination of BDE47 with resident time (1.5 years) in Australia for the single subject selected for half-life determination. Red dots shows actual measured concentration and blue lines shows median concentration and 2.5th and 97.5th percentiles of concentration estimates from 10,000 iterations (see Materials and methods for underlying assumptions). Green lines shows median and 2.5th and 97.5th percentiles of the Australian subjects.




Figure 2. Median estimate of intrinsic elimination half-life for tri- to hexaBDE congeners. Error bars indicates 2.5th and 97.5th percentile of Monte Carlo estimate.

References:

1. Thuresson K, Höglund P, Hagmar L et al. (2006) *Environ Health Perspect.* 114: 176-181.
2. Geyer HJ, Schramm K-W, Darnerud PO et al. (2002) *Organohalogen Comp.* 66: 3820-3825.
3. Wong F, Cousins IT, MacLeod M. (2013) *Environ Int.* 59: 168-174.
4. Sjödin A, Jones R, Lapeza C et al. (2004) *Anal. Chem.* 76: 1921-1927.
5. Jones R, Edenfield E, Anderson S et al. (2012) *Organohalogen Comp.* 74: 97-97.
6. Russell MH, Waterland RL, Wong F. (2015) *Chemosphere* 129: 210-216.
7. Harrison RL. (2010) *AIP Conf Proc.* 1204: 17-21.